

Mepilex® Border Sacrum and Mepilex® Border Heel

Designed to protect, proven at every layer

The problem

Pressure injuries (PIs) impact millions of patients annually, with:



12.8% of all adult patients

in hospitals developing a pressure injury¹.



8.4% of all hospital patients

acquiring a pressure injury during their stay¹.

Preventing PIs saves lives, protects patients and reduces the impact on the healthcare system. Our dressings play a crucial role in preventing and treating PIs.

The solution

The International Guidelines recommend the use of multilayer silicone foam dressings for pressure injury prevention (PIP)^{2,3}.



Mölnlycke dressings have the highest level of proof in PI prevention.

**19
17**

Randomised Controlled Trials (RCTs).

Meta-analyses or systematic reviews.



This commitment to high-quality research means that Mepilex® Border dressings

have more high-level evidence demonstrating their effectiveness in preventing pressure injuries^{4*}.

*Competitor dressings: Allevyn™ Life (Smith&Nephew), Aquacel® Foam (Convatec), Optifoam® Gentle (Medline), Biatain® Silicone (Coloplast)



Mepilex® Border Sacrum and Mepilex® Border Heel for prevention

Designed on four key pillars:

- Deep Defense® technology provides better protection and potentially longer wear time.⁵
- Up to 197% better conformability than other bordered foam dressings^{7,8}
- Unique structural integrity sustaining performance for up to 7 days.⁶
- Up to 65% wider surface area of protection versus competitors.⁹

Mepilex® Border Sacrum and Mepilex® Border Heel for treatment



Up to **74X faster** initial absorption, reducing risk of maceration¹¹



Up to **3.6X lower** swelling, keeping a discreet profile¹¹



Up to **130% higher fluid*** handling capacity than other leading foam dressings, supporting a potential longer wearing time^{12**}



Exudate progress monitoring, enabling undisturbed wound healing^{13,14}

↓ **64%**

In a US economic study within a cohort of 1.03M patients, Mepilex® Border

Sacrum demonstrated a 64% reduction in per-patient treatment cost compared to when no dressing was used¹⁰.

↓ **88%**

A landmark USA-based RCT demonstrated that using Mepilex® Border Sacrum as part of a

prevention protocol reduced the incidence of hospital-acquired sacral pressure injuries by up to 88% reduction in the incidence of PIs¹⁵.

Mepilex® Border Sacrum Dressing

Product code	Size	Pcs/box	Pcs/case	HCPCS
282055	6.3" x 7.9" (16 x 20 cm)	10	50	A6213
282455	8.7" x 9.8" (22 x 25 cm)	10	40	A6213

Mepilex® Border Heel Dressing

Product code	Size	Pcs/box	Pcs/case	HCPCS
282790	8.7" x 9.1" (22 x 23 cm)	10	10	A6210

References

1. Li Z, Lin F, Thalib L, et al. Global prevalence and incidence of pressure injuries in hospitalised adult patients: A systematic review and meta-analysis. International Journal of Nursing Studies. 2020;105:103546.
2. Association of periOperative Registered Nurses (AORN). (2022). Guideline for positioning the patient. In: Guidelines for Perioperative Practice. 3. European Pressure Ulcer Advisory Panel (EPUAP), National Pressure Injury Advisory Panel (NPIAP), and Pan Pacific Pressure Injury Alliance (PPPIA). Prevention and Treatment of Pressure Ulcers/Injuries: Clinical Practice Guideline. The International Guideline, 4th ed. Emily Haesler (Ed.). EPUAP/ NPIAP/PPPIA; 2025. 4. Mölnlycke Health Care. Data on File. 2020. 5. Levy, A., Gefen, A. Assessment of the biomechanical effects of prophylactic sacral dressings on tissue loads: A computational modeling analysis. Ostomy Wound Management 2017;63(10):48-55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29091038> [Accessed 4 February 2019]. Mepilex Border Sacrum. 6. Burton, J.N., Fredrickson, A.G., Capunay, C., Tanner, L., Oberg, C., Santamaria, N., Gefen, A., Call, E. Measuring tensile strength to better establish protective capacity of sacral prophylactic dressings over 7 days of laboratory aging. Advances in Skin and Wound Care 2019;32(7 Supplement):S21-S29. 7. Mölnlycke Health Care. Data on File. 2025. 8. Mölnlycke Health Care. Data on File. 2025. 9. Mölnlycke Health Care. Data on File. 2025. 10. Padula, W.V. Effectiveness and value of prophylactic 5-layer foam sacral dressings to prevent hospital-acquired pressure injuries in acute care hospitals: An observational cohort study. Journal of Wound, Ostomy, and Continence Nursing 2017 44(5):413-419. Available from: doi: 10.1097/WON.0000000000000358 11. Mölnlycke Health Care. Data on File. 2025. 12. Mölnlycke Health Care. Data on File. 2025. 13. Mölnlycke Health Care. Data on File. 2022. 14. Mölnlycke Health Care. Data on File. 2018. 15. Kalowes P, Messina V, Li M. Five-layered soft silicone foam dressing to prevent pressure ulcers in the intensive care unit. Am J Crit Care. 2016;25(6):e108-e119.

*Tested using a simulated fluid that mimics chronic wound exudate (SWF A). Data on file. **Other dressings test and compared with were Allevyn® Life Sacrum (Smith&Nephew), Optifoam Gentle SA Sacrum & Optifoam Gentle EX Sacrum (Medline), Urgotul Absorb Sacrum (Urgo), and Convafoam Silicone Sacrum (ConvaTec).

Mölnlycke Health Care AB, Entreprenörstråket 21, SE-431 21 Mölndal Sweden. Phone + 46 31 722 30 00. The Mölnlycke, Mepilex, Z-Flo and Safetac trademarks, names and logos are registered globally to one or more of the Mölnlycke Health Care group of companies. ©2025 Mölnlycke Health Care AB. All rights reserved. HQIM00830