Patient Risk Assessment



Clinician's name				
Patient ID				
Assessment date				
Type of surgery				
Tick boxes that a	pply			
Diabetes	Duration >120		ins	ASA≥3
Obesity	>75 th percentile (for procedure durations <120		ns <120 mins)	Wound class II (clean contaminated)
Risk (determined by ti	ick boxes above) —			· · · · · · · · · · · · · · · · · · ·
Low	Moderate	Elevated		High*
				⊘ ⊘ ⊘ +
Advanced Wound Dressing Product in Mepilex® Bor		ndication		Negative pressure wound therapy (ciNPWT)
				Product indication Avance® Solo
Risk profile Low	Moderate		Elevated	High
Recommended treatment				

www.molnlycke.com.au | www.molnlycke.co.nz

high regardless of the status of any other risk factor(s).

You will be required to input the following code **1FtYrV** or your **AHPRA registration number** to access our websites. Mölnlycke, 12 Narabang Way, Level 4, Belrose, NSW, 2085. Phone 1800 005 231. New Zealand Orders & Enquiries 0800 005 231, www.molnlycke.co.nz. The Mölnlycke, Avance and Mepilex trademarks, names and logos are registered globally to one or more of the Mölnlycke Health Care Group of Companies. © 2024 Mölnlycke Health Care. All rights reserved. AUWC_001101

* If the patient has wound class III or IV, or emergent surgery, the patient's risk status should be considered

