







Produkter til behandling av brannskader

Sårrens



Granudacyn®

Væskéhåndtering / Antimikrobiell lokalbehandling

Væskeniå	Områder som normalt er enkle å bandasjere	Områder som er vanskelige å bandasjere
Lavt	 Mepilex® Ag	 Mepilex® Ag
Medium	 Mepilex® Ag	 Mepilex® Transfer Ag + sekundærbandasje
Høyt	 Exufiber® Ag+ + sekundærbandasje	 Mepilex® Transfer Ag + sekundærbandasje

Sekundærbandasjer



Mextra®

Ved svært høyt væskeniå



Mesorb®

Ved høyt væskeniå

Fiksering



Tubifast®



Tubifast®
garments

Ødembehandling



Tubigrip®

Arrbehandling



Mepiform®

Ansvarsfraskrivelse: Les alltid bruksanvisningen som følger med det medisinske produktet før bruk. Utvalg og indikasjoner for bruk kan variere avhengig av land.

Referanser:

1. Upton D, Solowiej K. Pain and stress as contributors to delayed wound healing. *Wound Practice and Research* 2010, Vol. 18(3). 2. Gee Kee EL, Kimble RM, Cuttle L, Khan A, Stockton KA. Randomized controlled trial of three burns dressings for partial thickness burns in children. *Burns* 2015, 41(5):946-955. 3. Silverstein P, Heimbach D, Meltes H et al. An open, parallel, randomized, comparative, multicenter study to evaluate the cost-effectiveness, performance, tolerance, and safety of a silver containing soft silicone foam dressing (intervention) vs silver sulfadiazine cream. *J Burn Care Res* 2011, 32(6): 617-626. 4. Gee Kee EL, Stockton K, Kimble RM et al. Cost-effectiveness of silver dressings for paediatric partial thickness burns. An economic evaluation from a randomized controlled trial. *Burns* 2017, 43(4): 724-732. 5. Aggarwala S, Harish V, Roberts S et al. Treatment of partial thickness burns: a prospective, randomised controlled trial comparing Biobrane, Acticoat, Mepilex Ag and Aquacel Ag. *J Burn Care Res* 2020, 42(5): 934-43. 6. Tang H, Lv G, Fu J et al. An open, parallel, randomized, comparative, multicenter investigation evaluating the efficacy and tolerability of Mepilex Ag versus silver sulfadiazine in the treatment of deep partial-thickness burn injuries. *J Trauma Acute Care Surg* 2015, 78(5): 1000-1007. 7. Gleeson AP, Stuart MJ, Wilson B, Phillips B. Ultrasound assessment and conservative management of inversion injuries of the ankle in children: Plaster of Paris versus tubigrip. *Journal of Bone and Joint Surgery - Series B* 1996, 78(3):484-7. 8. Naem M, Rahimnadjad MK, Rahimnadjad NA, Idrees Z, Shah GA, Abbas G. Assessment of functional treatment versus plaster of Paris in the treatment of grade 1 and 2 lateral ankle sprains. *Journal of Orthopaedics and Traumatology* 2014, 16(1):41-6. 9. Mölnlycke Health Care. Data on file. Tubigrip. 10. Eytier C, Gazeau E, Beneteau G, Verfaille G. Convenience and tolerance of the combination of a soft silicone foam dressing and a two-way stretch tubular bandage in the management of local wounds. *Journal des plaies et cicatrisations* 2013, 18(88):38-44. 11. Mölnlycke Health Care. Data on file. Tubifast. 12. Mölnlycke Health Care. Data on File. Mepiform. 13. Wigger-Albert W, Kuhlmann M, Wilhelm D, Mrowietz U, Eichhorn K, Ortega J, et al. Efficacy of a polyurethane dressing versus a soft silicone sheet on hypertrophic scars. *Journal of wound care* 2009, 18(5):208. 10-4. 14. Armstrong DG, Bohn G, Glat P, Kavros SJ, Kirsner R, Snyder R et al. Expert recommendations for the use of Hypochlorous solution: science and clinical application. *Ostomy Wound Manage* 2015, 61 (5 suppl): 45-18S 15. Consensus on Wound Antisepsis: Update 2018. *Skin Pharmacol Physiol* 2018;31:28-58. 16. Chadwick P, Taherinejad F, Hamberg K, Waring M. Clinical and scientific data on a silver-containing soft-silicone foam dressing: an overview. *Journal of Wound Care* 2009, 18(11):483-491. 17. Mölnlycke Health Care. Data on file. Mepilex Transfer Ag. 18. Schweiger H, Smith D, Cruse CW et al. An open, non-controlled, single-centre, clinical investigation to evaluate efficacy when using a soft silicone wound contact layer containing silver. Poster presentation at the 15th European Burns Association Congress, Vienna, Austria, 2013. 19. Chadwick P, McCardle J. Open, non-comparative, multicenter post clinical study of the performance and safety of a gelling fibre wound dressing on diabetic foot ulcers. *Journal of Wound Care* 2016, 25(4): 290-300. 20. Smet, S., Beele, H., Saine, L., Suys, E., Henrickx, B. Open, non-comparative, multi-centre post market clinical follow-up investigation to evaluate performance and safety on pressure ulcers when using a gelling fibre dressing as intended. Poster Presentation at European Pressure Ulcer Advisory Panel Conference 2015, Ghent, Belgium. 21. Mölnlycke Health Care. Data on file. Exufiber. 22. David F, et al. A randomised, controlled, non-inferiority trial comparing the performance of a soft silicone-coated wound contact layer (Mepitel One) with a lipidocolloid wound contact layer (UrgoTul) in the treatment of acute wounds. *International Wound Journal* 2018, 23. Mölnlycke Health Care. Data on file. Avance Solo. 24. Nischwitz SP, Luze H, Popp D, Winter R, Draschl A, Schellnegger M, Kargl L, Rappl T, Giretzlehner M, Kamolz LP. Global burn care and the ideal burn dressing reloaded—A survey of global experts. *Burns* 2021, 47: 1665-1674.

Les mer på www.molnlycke.no

Mölnlycke Health Care AS, postboks 6229, Etterstad, 0603 Oslo. Tlf: 22 70 63 70. The Mölnlycke, Mepilex, Mextra, Tubifast, Exufiber, Mepitel, Mepiform, Mesorb, Granudacyn, Tubigrip, Melgisorb, Avance and Safetac trademarks, names and logotypes are registered globally to one or more of the Mölnlycke Health Care group of companies.
© 2025 Mölnlycke Health Care AB. All rights reserved. 2023NOWC3742503



Mölnlycke®